

Supporting Information

Supramolecular assembly of Toll-like receptor 7/8 agonist into multimeric water-soluble constructs enables superior immune stimulation in vitro and in vivo

Alexander K. Andrianov^{,†}, Alexander Marin[†], Ruixue Wang[†], Hatice Karauzum[#], Ananda Chowdhury[†], Pragati Agnihotri^{§,o}, Abdul S. Yunus[†], Roy A. Mariuzza^{†,§,o}, Thomas R. Fuerst^{†,§}*

[†]Institute for Bioscience and Biotechnology Research, University of Maryland, Rockville, Maryland, 20850, USA.

[#]Integrated BioTherapeutics, Inc., Rockville, MD 20850, USA.

[§]Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, Maryland, 20742, USA.

^oW.M. Keck Laboratory for Structural Biology, University of Maryland Institute for Bioscience and Biotechnology Research, Rockville, Maryland 20850

Corresponding author

Alexander K. Andrianov: *E-mail: aandrianov@umd.edu

Table of Content


Scheme S1. Chemical reaction scheme of PCPP and RSQD S-2

Scheme S2. Experimental microfluidics set-up for preparing PCPP-R formulations S-3


Figure S1. ^1H NMR spectrum of PCPP-RSQD formulation S-4

Figure S2. DLS profiles of PCPP-R – E2, PCPP-R and PCPP S-5

Figure S3. Cell mediated responses.....S-6

Scheme S1. Chemical reaction scheme of PCPP and RSQD.

Scheme S2. Experimental microfluidics set-up for preparing of PCPP-R formulations.






Figure S1. ¹H NMR spectrum of PCPP-RSQD formulation (RSQD:COO⁻(PCPP)=1:1 (mol/equiv), 1M KOH in D₂O; peak assignments are in Fig. 2E).




Figure S2. DLS profiles of PCPP-R – E2, PCPP-R and PCPP (0.125 mg/mL E2, 0.125 mg/mL PCPP, 0.062 mg/mL RSQD, PBS, pH 7.4).




Figure S3. Cell mediated responses to E2 peptides as assessed ex-vivo in splenocytes derived from immunized BALB/c mice with various formulations: (A) representative FACS plots and (B) summary data of TNF- α single (Y axis), IFN- γ single (X axis) and TNF- α /IFN- γ double producing cells (top right quadrant).